

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**



Eur päisches Patentamt  
Eur pean Patent Office  
Offic eur péen des br vets



(11)

**EP 0 727 430 A1**

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:  
21.08.1996 Bulletin 1996/34

(51) Int Cl.<sup>6</sup>: C07F 15/00, A61K 31/28

(21) Application number: 96300601.0

(22) Date of filing: 29.01.1996

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT  
SE

(30) Priority: 14.02.1995 GB 9502799



(71) Applicant: JOHNSON MATTHEY PUBLIC LIMITED  
COMPANY  
London EC1N 8JP (GB)

(72) Inventor: Murrer, Barry A.  
Reading RG1 5SB, Berkshire (GB)

(74) Representative: Wishart, Ian Carmichael et al  
Patents Department  
Johnson Matthey Technology Centre  
Blounts Court  
Sonning Common  
Reading, Berks RG4 9NH (GB)

(54) Improvements in PT complexes

(57) Platinum complexes of formula Ia or Ib



where

each A, is a leaving group and may be the same or different, which is suitably halo, especially chloro, hydroxy, carboxylate or together form a bi-dentate carboxylate or sulphate,

each B, which may be the same or different, is halo, hydroxy, carboxylate, carbamate or carbonate ester,

Z is a substituted amine wherein the substituent sterically hinders access of the Pt atom to a DNA strand of a tumour cell, and

X is NH<sub>3</sub> or mono- or di-alkyl substituted NH<sub>3</sub>, are active against cancer cells, and appear to show a unique form of chemical and biological action.


EP 0 727 430 A1

## Description

The present invention concerns improvements in platinum complexes, and particularly concerns novel platinum complexes having activity against cancer cells.

The activity of a Pt(II) complex, cisplatin,  $[\text{PtCl}_2(\text{NH}_3)_2]$ , against cancer cells was discovered some twenty years ago, and this complex has become a major pharmaceutical. However, cisplatin has long been known to suffer from two major drawbacks; namely, its severe toxicities (especially nephrotoxicity, nausea and vomiting and neurotoxicity) and the propensity of many tumours to exhibit resistance (either intrinsically or acquired after an initial promising response). Many platinum complexes have been studied in an attempt to overcome the limitations of cisplatin.

The present invention provides a cis-platinum complex of general formula Ia or Ib



where

each A, is a leaving group and may be the same or different, which is suitably halo, especially chloro, hydroxy, carboxylate or together form a bi-dentate carboxylate or sulphate,

each B, which may be the same or different, is halo, hydroxy, carboxylate, carbamate or carbonate ester,

Z is a substituted amine wherein the substituent sterically hinders access of the Pt atom to a DNA strand of a tumour cell, and

X is  $\text{NH}_3$  or mono- or di-alkyl substituted  $\text{NH}_3$ .

Preferably the complex is of formula Ia.

Particularly, we have found that where Z is an unsaturated cyclic amine, where the ring may contain one or more other hetero-atoms, and most especially where Z is pyridine, and where Z has a substituent on the atom adjacent to the amine nitrogen atom, the complex appears to be sterically hindered.

The cyclic amine may be a 5- or 6-membered monocyclic or 8 to 10-membered polycyclic amine, especially bicyclic, for example fused ring systems where the amine is co-ordinated through the nitrogen atom of a pyridine ring. In the case of such bicyclic fused ring systems, the other ring may be phenylene or may contain one or more heteroatoms, especially nitrogen or oxygen.

In the case of substituted cyclic amines, the substituent may be lower alkyl or alkoxy of 1 to 4 carbon atoms, (especially methyl or methoxy), halo, (especially chloro or bromo), or aryl, (especially benzyl). The substituent may itself be substituted by lower alkyl or halo. By the term "lower alkyl" is meant an alkyl group with from 1 to 6 carbon atoms. The cyclic amine may carry other substituents either adjacent to the coordinating nitrogen atom or elsewhere on the ring. Other substituents include electron-withdrawing or electron-donating substituents such as nitro and alkoxy *eg* methoxy. If the cyclic amine is a fused ring system where the fused ring is an aromatic ring in positions 2 and 3 of the cyclic amine, no other substituent is necessary, although a substituent may be present.

The leaving groups A may be selected from those well known to those skilled in the art.

Preferably, each A is the same, and is chloro, or together form cyclobutane-1,1-dicarboxylate or sulphate. In the case of Pt(IV) complexes of formula Ib, preferably each B is the same, and preferably is hydroxy.

We have demonstrated that members of this novel class of sterically hindered complexes exhibit different chemical characteristics from analogues which are not hindered, and they exhibit a different biological profile. Tests relating to these characteristics are described below.

The complexes of formula Ia and Ib are novel and may be prepared by methods analogous to those described in the art. For example, general preparations of platinum complexes with mixed ammine/alkylamine ligands are given by P D Braddock *et al* Chem. Biol. Interactions 1975, II, 145. We refer also to EP 328274 (Johnson Matthey).

The requisite starting materials are themselves known.

The complexes of the invention demonstrate activity against cancer cells *in vitro*; some complexes have also been tested *in vivo* and have also demonstrated interesting activity. The complexes appear to show a different activity, particularly towards cancer cells which are resistant to treatment with cisplatin, to the existing commercial platinum anti-cancer complexes and therefore the complexes are indicated for use in the treatment of tumours.

According to a further aspect of the invention, there is provided a complex of formula Ia or Ib for use in medicine

and in particular for use in the treatment of cancer. A still further aspect of the invention provides the use of a complex of formula Ia or Ib in the manufacture of a medicament for the treatment of cancer. Alternatively, there is provided a method of treating cancer comprising administering to a patient in need thereof a pharmaceutically effective amount of a complex of formula Ia or Ib.

The active complexes may be administered in the form of pharmaceutical compositions formulated according to well known principles. Thus, a still further aspect of the present invention provides a pharmaceutical composition comprising a compound of formula Ia or Ib in admixture with a pharmaceutically acceptable diluent or carrier and optionally one or more other therapeutic agents. Such compositions may be in the form of solutions or suspensions for injection, or to be in capsule, tablet, dragee, or other solid composition or as a solution or suspension for oral administration, or formulated into pessaries or suppositories, or sustained release form of any of the above. Suitable diluents, carriers, excipients and other components are known. It may be desirable also to formulate a composition for topical administration such as an ointment or cream, or to be administered as a transdermal patch.

The pharmaceutical compositions according to the invention may contain dosages determined in accordance with conventional pharmaceutical methods, suitably to provide active compounds in the dosage range in humans of from 0.1 to 100mg/kg body weight per day, in a single unit dose or in a number of smaller unit doses. Preferred dosage ranges are 1 to 30mg/kg body weight per day.

The complexes of the invention may be administered alone or in combination with another chemotherapeutic agent, such as cisplatin, either as a single treatment or course of treatment or as part of combined therapy with other pharmaceuticals to overcome or diminish side effects or to improve bio-availability, or in combination with other therapies such as radiation treatment.

Without wishing to be bound by any theory, it appears to the inventors that the complexes of the invention exhibit increased selectivity towards DNA in cancer cells compared to unhindered analogues, especially cisplatin. It has been reported by Tobe *et al* (Wadley Medical Bulletin 7, 1, 114-135) that steric effects of bulky substituents close to the platinum atom have a profound effect upon the rate of substitution reactions there. At that time no systematic study of the way in which variation of the substituent on the primary amine affects the rate and extent of bonding of the Pt complex to DNA had been undertaken. There is no evidence as to whether such effects could be beneficial or detrimental to the antitumour activity of platinum complexes. The present invention introduces steric hindrance as close as possible to the platinum atom by the use of substituted heterocyclic ligands, rather than amines, and we believe that the complexes of the invention show unexpected benefits. In particular, the complexes of the invention exhibit reduced resistance factors (resistance factor is the ratio of the activity of a complex against a parent line of cancer cells to the activity against a derivative of that cell line which has developed resistance to cisplatin). For example, in the CH1 xenograft (Institute of Cancer Research, Sutton, Surrey) which is derived from cell line sensitive to cisplatin, cisplatin shows significant activity at the maximum tolerated dose, however the tumour does grow back. The novel complex  $[PtCl_2(NH_3)(2\text{-methylpyridine})]$  shows slightly less potency, meaning that a greater dose is required than cisplatin, but the complex exhibits a curative effect, and the tumour does not grow back.

We believe that cisplatin, and all known anti-tumour active Pt complexes, bind to DNA in cancer cells, causing cross-linking of DNA strands. Approximately 10% of the total links are interstrand cross-links. Our tests with certain of the complexes of the invention indicate that there is no detectable cross-linking between strands. This indicates a different mechanism of action.

The invention will now be described with reference to the following preparative examples.

#### EXAMPLE 1

##### (SP-4-3)-amminedichloro(2-methylpyridine)platinum(II)

KCl (2.2g, 29.5mmol) and  $K[PtCl_3(NH_3)]$  (10g, 28mmol) were dissolved in water (100ml) and stirred during the addition of KI (13.90g, 84mmol) in water (15ml). 2-methylpyridine (2.8g, 30mmol) was then added. After 3 hours stirring, the yellow precipitate of (SP-4-3)-amminediiodo(2-methylpyridine)platinum(II) was collected by filtration, washed with water and then methanol and dried *in vacuo*, yield 8.06g. The solid was added to a stirred solution of  $AgNO_3$  (4.78g, 28.1mmol) in water (50ml) and stirring continued for 6 hours in the dark. After filtration to remove silver halides, the filtrate was treated with concentrated HCl (4.5ml, ca 49.5mmol) and stirred for 3 days. The resulting pale yellow solid was collected by filtration, washed with water then acetone, and dried *in vacuo* to give (SP-4-3)-amminedichloro(2-methylpyridine)platinum(II) (5.78g, 52%). Found: C 19.4, H 2.7, N 7.3, Cl 18.9%;  $C_6H_{10}N_2Cl_2Pt$  requires C 19.5, H 2.7, N 7.4, Cl 18.9%.

Using analogous procedures, the following compounds were prepared:

	Ligand Z	% yield	Microanalysis % found (expected)			
			C	H	N	Cl
Example 2	2-ethylpyridine	41	21.5 (21.5)	3.2 (3.1)	7.3 (7.2)	18.0 (18.2)
Example 3	2-methylquinolin	17	28.4 (28.0)	3.0 (2.8)	6.5 (6.5)	16.6 (16.6)
Example 4	2,4-dimethylpyridine	37	21.4 (21.5)	3.2 (3.1)	7.1 (7.2)	18.1 (18.2)
Example 5	2-isopropylpyridine	48	23.9 (23.8)	3.6 (3.5)	6.8 (6.9)	17.3 (17.6)
Example 6	2,6-dimethylpyridine	14	21.5 (21.5)	2.85 (3.1)	7.3 (7.2)	18.4 (18.2)

**EXAMPLE 7****(OC-6-43)-amminedichlorodihydroxo(2,4-dimethylpyridine)platinum(IV)**

The product of Example 4, [(SP-4-3)-amminedichloro(2,4-dimethylpyridine)platinum(II)], (2.0g), heptane (5ml), water (2.9ml) and H<sub>2</sub>O<sub>2</sub> (30% w/v 2.9ml) were stirred vigorously and refluxed for 2 hours. On cooling, the yellow solid was collected by filtration, washed with H<sub>2</sub>O and dried to give (OC-6-43)-amminedichlorodihydroxo(2,4-dimethylpyridine)platinum(IV) (1.74g, 77%). Found C 19.77, H 3.03, N 6.52, Cl 16.57%, C<sub>7</sub>H<sub>14</sub>N<sub>2</sub>Cl<sub>2</sub>O<sub>2</sub>Pt requires C 19.81, H 3.30, N 6.60, Cl 16.75%.

**EXAMPLE 8****(OC-6-43)-amminebisbutyratodichloro(2,4-dimethylpyridine)platinum(IV)**

The product of Example 7 (OC-6-43)-amminedichlorodihydroxo(2,4-dimethylpyridine)platinum(IV) (1.5g) was suspended in butyric anhydride (5.6g) and stirred at room temperature for 3 days. The pale yellow solid was collected by filtration, washed thoroughly with MeOH and dried *in vacuo* to give (OC-6-43)-amminebisbutyratodichloro(2,4-dimethylpyridine)platinum(IV) (1.17g). Found C 31.4, H 4.5, N 4.9, Cl 12.5; C<sub>15</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>Cl<sub>2</sub>Pt requires C 31.9, H 4.6, N 5.0, Cl 12.6%.

**EXAMPLE 9****ammine(cyclobutane-1,1-dicarboxylato)(2,4-dimethylpyridine)platinum(II)**

2,4-dimethylpyridine (1.65g) was added to a solution of K[PtCl<sub>3</sub>(NH<sub>3</sub>)] (5.0g), KCl (1.1g) and KI (6.95g) in water (60ml). After 3 hours stirring at room temperature the yellow precipitate was collected by filtration, washed with water and then methanol and dried *in vacuo*. This compound was added to a solution of AgNO<sub>3</sub> (2.94g, 1.95mmol) in water (30ml) and acetone (5ml) and stirred in the dark for 5 hours. The mixture was filtered to remove silver halides and the filtrate was added dropwise to a warm (60°C) solution of dipotassium cyclobutane-1,1-dicarboxylate (13.2mmol). The solution was kept at 60°C for 2 hours, then left at room temperature overnight. The solvent was removed *in vacuo* and the white solid dissolved in the minimum amount of warm EtOH, filtered and stored at -20°C. After 4 hours a white crystalline material was filtered off, washed with cold ethanol and dried (yield 0.81g). NMR indicated the desired compound, together with ca 0.9mol EtOH/Pt. (NMR, CD<sub>3</sub>OD δ 8.7 d (1H), 7.3 s (1H), 7.1 d (1H), 3.6 q (CH<sub>2</sub>CH<sub>2</sub>OH), 3.1 s (3H), 2.9 m (4H), 2.4 s (3H), 1.9 m (2H), 1.2 t (CH<sub>3</sub>CH<sub>2</sub>OH). Microanalysis found C 34.8, H 5.1, N 5.3, C<sub>13</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>Pt. 0.9 EtOH requires C 35.3, H 4.7, N 5.6%, IR 1629cm<sup>-1</sup> (bound carboxylate).

The complexes of the invention were tested against human cancer cell lines grown in cell culture according to established testing procedures (described by Kelland *et al*, Cancer Research 53, 2581-2586, June 1993). The results are shown below in Table 1, in comparison with the commercial complexes cisplatin and carboplatin. The results are given in concentrations (μM) necessary to achieve a 50% decrease in cell proliferation, with resistance factors in brackets for the cisplatin-resistant cell lines. HX62 and SKOV3 are inherently resistant to cisplatin, and the cell lines bred to be resistant to cisplatin have the designation R, as in 41MR. Resistance factors for HX62 and SKOV3, for which there is no sensitive counterpart, have been derived by dividing by the IC<sub>50</sub> for the 41M line, an inherently cisplatin sensitive line.

It is not, of course, surprising that the resistance factors for cisplatin itself are fairly high, but it can be seen that the other commercial complex, carboplatin, is also less effective with cisplatin-resistant cell lines. The complexes of the invention appear to show reduced resistance factors compared to the commercial substances, and this is assessed

as being advantageous.

TABLE I

Compound	HX62	SKOV3	41M	41MR	CHI	CHIR	A2780	A2780R
Cisplatin	15.5 (37.8)	4.3 (10.5)	0.41	1.9 (4.6)	0.12	0.44 (3.7)	0.35	7.8 (22)
Carboplatin	85 (26)	37.7 (11)	3.3	8.8 (2.7)	1.3	6 (4.6)	2.6	37.4 (14)
Example 1	40 (9.8)	17.5 (4.3)	4.1	5 (1.2)	1.9	3.7 (1.9)	1.28	4.4 (3.4)
Example 2	36 (10)	20 (2.9)	3.4	5.7 (1.7)	1.4	3.6 (2.6)	4.4	11.0 (2.5)
Example 4	4.9 (1)	14.0 (3)	4.25	1.9 (0.4)	2.3	5.3 (2.3)	6.6	18.0 (2.7)
Example 5	49 (8)	20 (3.3)	6.1	5.6 (0.9)	2.4	4.2 (1.7)	3.9	14 (3.6)
Example 6	62 (2)	46 (1.4)	33	28 (0.84)	20	35 (1.74)	35	77 (2.2)
Example 7	72 (9)	47 (5.9)	8	17.5 (2.2)	3.1	8.9 (2.5)	14	30 (2.1)

The compound of Example 1 was used for toxicological studies and pharmacological studies in Balb C mice. The results are given in Table 2.

TABLE 2


TUMOUR	EXAMPLE I			CISPLATIN		
	LD <sub>50</sub>	ED <sub>90</sub>	TI	LD <sub>50</sub>	ED <sub>90</sub>	TI
<u>ADJ/PC6</u>						
Single dose (i.p.)	43	3	14.3	11.3	1.6	7.1
Single dose (p.o.)	560	6.2	90.3	140	24	5.8
Daily x 5 (i.p.)	21	0.8	26.3	2.4	0.2	11.7

LD<sub>50</sub> (mg/kg); dose causing 50% lethalityED<sub>90</sub> (mg/kg); dose required to reduce tumour mass by 90%TI, therapeutic index; LD<sub>50</sub>/ED<sub>90</sub>.

No significant histological evidence of liver, kidney, spleen, gastrointestinal tract, brain or skin toxicity was observed in mice receiving a LD<sub>10</sub> dose of the compound of Example 1. There were no significant differences in levels of alkaline phosphatase (ALP), alanine aminotransferase (ALT), the gut disaccharides, maltose, sucrose and trehalose, and urea and creatinine in treated mice (LD<sub>50</sub> dose) compared to control mice.

### Claims

1. A cis-platinum complex of general formula Ia or Ib



where

each A, is a leaving group and may be the same or different, or together form a bi-dentate carboxylate or sulphate,

each B, which may be the same or different, is halo, hydroxy, carboxylate, carbamate or carbonate ester,

Z is a substituted amine wherein the substituent sterically hinders access of the Pt atom to a DNA strand of a tumour cell, and

X is NH<sub>3</sub> or mono- or di-alkyl substituted NH<sub>3</sub>.

2. A complex according to claim 1, wherein Z is a 5- or 6- membered monocyclic or a 8 to 10-membered polycyclic amine.
3. A complex according to claim 2, wherein Z is a substituted pyridine or a bicyclic amine where the amine is coordinated through the nitrogen atom of a pyridine ring.
4. A complex according to claim 3, wherein Z is a pyridine substituted by lower alkyl or alkoxy in the 2 position.
5. A complex according to any one of the preceding claims, wherein each A is halo, hydroxy, carboxylate or together form bi-dentate carboxylate or sulphate.
6. A complex according to claim 1 which is selected from  
 (SP-4-3)-amminedichloro(2-methylpyridine) platinum(II); (SP-4-3)-amminedichloro(2-ethylpyridine) platinum(II); (SP-4-3)-amminedichloro(2-methylquinoline) platinum(II); (SP-4-3)-amminedichloro(2,4-dimethylpyridine) platinum(II); (SP-4-3)-amminedichloro(2-isopropylpyridine) platinum(II); (SP-4-3)-amminedichloro



(2,6-dimethylpyridine) platinum(II); (OC-6-43)-ammine dichlorodihydroxo-(2,4-dimethylpyridine) platinum(IV); (OC-6-43)-amminebisbutyratodichloro(2,4-dimethylpyridine) platinum(IV); and ammine(cyclobutane-1,1-dicarboxylato)(2,4-dimethylpyridine) platinum(II).

- 5 7. A pharmaceutical composition comprising as active ingredient a complex according to any one of the preceding claims, in admixture with a pharmaceutically acceptable diluent or carrier and optionally one or more other therapeutic agents.
- 10 8. A composition according to claim 7, in unit dosage form.
9. A composition according to claim 7 or 8 for oral administration.
10. A complex according to any one of claims 1 to 6 for use in medicine.
- 15 11. The use of a complex according to any one of claims 1 to 6 in the manufacture of a medicament for the treatment of cancer.

20

25

30

35

40

45

50

55



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 96 30 0601

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	CHEMICAL ABSTRACTS, vol. 121, no. 26, 26 December 1994 Columbus, Ohio, US; abstract no. 314427d, DUPONT, J. ET AL.: "CIS AND TRANS NUCLEOPHILIC ADDITIONS ON C...C BONDS ASSISTED BY PT(II) COMPLEXES. X-RAY CRYSTAL STRUCTURE OF A TRANS PT COMPOUND" XP002004297 * abstract * & POLYHEDRON, vol. 13, no. 17, 1994, pages 2583-2587,	1	C07F15/00 A61K31/28
X	CHEMICAL ABSTRACTS, vol. 103, no. 26, 30 December 1985 Columbus, Ohio, US; abstract no. 226258, POINTEAU, P. ET AL.: "SYNTHESIS AND STRUCTURE OF PLATINUM(II) (AMINOPHENOL) (PYRIDINE) COMPLEXES" XP002004298 * abstract * & EUR. J. MED. CHEM.--CHIM. THER., vol. 20, no. 4, 1995, pages 327-332,	1	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07F A61K
A	EP-A-0 199 524 (ENGELHARD CORPORATION) * the whole document * --- -/--	1	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 30 May 1996	Examiner Rinkel, L
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 150 (01.92) (P4/C01)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 96 30 0601

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.6)
A	JOURNAL OF MEDICINAL CHEMISTRY, 1989, pages 128-136, XP002004296 HOLLIS, L.S. ET AL.: "CHEMICAL AND BIOLOGICAL PROPERTIES OF A NEW SERIES OF CIS-DIAMMINEPLATINUM(II) ANTITUMOR AGENTS CONTAINING THREE NITROGEN DONORS: CIS-(PT(NH3)2(N-DONOR)CL)+" * the whole document *	1	TECHNICAL FIELDS SEARCHED (Int. CL.6)
A	EP-A-0 328 274 (JOHNSON MATTHEY, INC.) * the whole document *	1	
A	EP-A-0 503 830 (JOHNSON MATTHEY PLC) * the whole document *	1	
A	EP-A-0 333 351 (JOHNSON MATTHEY PLC) * the whole document *	1	
A	EP-A-0 273 315 (SHIONOGI & CO. LTD) * the whole document *	1	
A	EP-A-0 115 929 (TANABE SEIYAKU CO., LTD) * the whole document *	1	
P,X	WO-A-95 07698 (THE UNIVERSITY OF VERMONT AND STATE AGRICULTURAL COLLEGE) 23 March 1995 * examples 2,5,9 *	1	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 30 May 1996	Examiner Rinkel, L
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 150 (01/92) (P/0401)

